

A NEW PROOF FOR THE MULTIPLICATIVE PROPERTY OF THE BOOLEAN CUMULANTS WITH APPLICATIONS TO OPERATOR-VALUED CASE

MIHAI POPA

ABSTRACT. The paper presents several combinatorial properties of the boolean cumulants. A corollary is a new proof of the multiplicative property of the boolean cumulant series that can be easily adapted for the case of boolean independence with amalgamation over an algebra.

AMS Subject Classification: 45L53, 46L08

Keywords: boolean cumulants, multilinear function series

1. INTRODUCTION

Boolean probability theory, according to [11] is one of the three symmetric universal probability theories (the other two being classical and free probability theories). It have been in the literature at least since early '70's ([15]) with various developments, from stochastic differential equations ([14]) to measure theory ([13], [2]) and Appell polynomials ([1]).

Two important notions in free probability theory are the R and S transforms. For X a non-commutative random variable of finite moments, $R_X(z)$ and $S_X(z)$ are power series that encode the information from the moment generating function of X and have the following additive, respectively multiplicative, property: if X, Y are free, then $R_{X+Y} = R_X = R_Y$ and $S_{XY} = S_X S_Y$.

The notion of boolean cumulants has been in the literature, in various contexts, at least since early '70's ([15], [14], [8]). Their generating function, that we will call the B -transform, has an additive property similar to the R -transform ([13]): if X and Y are boolean independent, then $B_{X+Y} = B_X + B_Y$.

In [6], U. Franz remarked an interesting multiplicative property of the B -transform, which gives a valuable tool for studying the multiplicative boolean convolution. For X, Y boolean independent, one has that

$$(1) \quad B_{(1+X)(1+Y)} = B_{(1+X)} B_{(1+Y)}.$$

In literature there are two proves of 1. The original proof, in [6], uses algebraic properties of the resolvent function. The proof in [2], in the spirit of D. Voiculescu's proof of the property for the R and S -transforms, uses a Fock-type model and properties of the Cauchy transform. In the attempt to develop a boolean probability theory with amalgamation over an algebra, when scalar-valued functionals are replaced by conditional expectations, the above approaches are not suitable to prove a similar multiplicative property, since objects such as the Cauchy transform and resolvent function do not have natural analogues. The present paper gives a combinatorial proof of (1), which is very easily adapted for the amalgamation over an algebra case (see Section 4).

In [4], [5], K. Dykema develops an analogue for the S -transform in the operator-valued free probability theory and remarks that, due to non-commutativity, it satisfy a “twisted” multiplicative property. Namely, if $S = T^{-1}$, then $T_{XY} = T_X(T_Y I T_Y^{-1})T_Y$. Corollary 4.6 from Section 4 shows that the operator-valued B -transform has the same multiplicative property as in the scalar case, without any “twists”.

2. PRELIMINARIES

Let $\mathcal{A}$ and $\mathcal{C}$ be a unital algebras and $\varphi : \mathcal{A} \longrightarrow \mathcal{C}$ be a linear functional with $\varphi(1) = 1$. The typical situation is when $\mathcal{C}$ is $\mathbb{C}$ or a subalgebra of $\mathcal{A}$, but we require it, nor that $\mathcal{C}$ is commutative.

The subalgebras $\mathcal{A}_1$ and $\mathcal{A}_2$ of $\mathcal{A}$ are said to be boolean independent with respect to φ (or just boolean independent) if

$$(2) \quad \varphi(X_1 Y_1 X_2 \cdots) = \varphi(X_1) \varphi(Y_1) \cdots$$

for all $X_1, X_2, \dots \in \mathcal{A}_1$ and $Y_1, Y_2, \dots \in \mathcal{A}_2$.

If $\mathcal{A}_1$ and $\mathcal{A}_2$ are subalgebras of $\mathcal{A}$, we will denote by $\mathcal{A}_1 \vee \mathcal{A}_2$ the algebra they generate in $\mathcal{A}$. With this notation, 2 can be interpreted as follows: if $\mathcal{A}_1$ and $\mathcal{A}_2$ are two boolean independent subalgebras of $\mathcal{A}$, then

$$(3) \quad \varphi(a_1 X Y a_2) = \varphi(a_1 X) \varphi(Y a_2)$$

for all $X \in \mathcal{A}_1, Y \in \mathcal{A}_2$ and $a_1, a_2 \in \mathcal{A}_1 \vee \mathcal{A}_2$.

We define the boolean cumulant of order n as the multilinear function $b^n : \mathcal{A}^n \longrightarrow \mathcal{C}$ given by the recurrence

$$(4) \quad \varphi(a_1 \cdots a_n) = \sum_{k=1}^n b^k(a_1, \dots, a_k) \varphi(a_{k+1} \cdots a_n).$$

Note that (4) is equivalent to

$$\begin{aligned} \varphi(a_1 \cdots a_n) &= \sum_{k=1}^n \varphi(a_1, \dots, a_k) b^{n-k}(a_{k+1} \cdots a_n) \\ &= \sum b^{k_1}(a_1, \dots, a_{k_1}) b^{k_2-k_1}(a_{k_1+1}, \dots, a_{k_2}) \cdots b^{n-k_m}(a_{k_m+1}, \dots, a_n) \end{aligned}$$

where the last summation is done over all $0 \leq m \leq n-1$ and all the m -uples $1 \leq k_1 < \cdots < k_m$. We will write b_X^n for $b^n(X, \dots, X)$. For the moment, respectively boolean cumulant generating power series of X we will use the notations $M_X(z)$, respectively $B_X(z)$, i.e.

$$\begin{aligned} M_X(z) &= \sum_{n=1}^{\infty} \varphi(X^n) z^{n-1} \\ B_X(z) &= \sum_{n=1}^{\infty} b_X^n z^{n-1}. \end{aligned}$$

The computations in the following two sections will often involve functions of many arguments. For brevity sake, we will use the short-hand notation described below:

If I is a set of indices, $\{a_i\}_{i \in I}$ is a family of elements from $\mathcal{A}$ and $\vec{u} = (i_1, \dots, i_n)$ is an ordered sequence from I , we will write $\varphi(a_{\vec{u}})$ for $\varphi(a_{i_1} a_{i_2} \cdots a_{i_n})$, respectively $b^n(a_{\vec{u}})$ for $b^n(a_{i_1}, a_{i_2}, \dots, a_{i_n})$. The length (cardinality) of $\vec{u}$ will be denoted by

$|\vec{u}|$. If $\vec{u}_1, \vec{u}_2$ are two ordered sequences from I , their ordered concatenation will be denoted by $\overrightarrow{u_1, u_2}$.

3. PROPERTIES OF THE BOOLEAN CUMULANTS

The following property is analogue to the vanishing of free cumulant with free independent entries. For the boolean case, only a weaker property is true.

Proposition 3.1. *Let $\mathcal{A}_1, \mathcal{A}_2$ be boolean independent subalgebras of $\mathcal{A}$. If $n, m \geq 0$ and $a_1, \dots, a_{n+m+2} \in \mathcal{A}_1 \vee \mathcal{A}_2$ such that $a_{n+1} = X \in \mathcal{A}_1$, $a_{n+2} = Y \in \mathcal{A}_2$, then:*

$$b^{n+m+2}(a_1, \dots, a_n, X, Y, a_{n+3}, \dots, a_{n+m+2}) = 0$$

Proof. Let $\vec{w} = (1, 2, \dots, n+m+2)$, $\vec{u} = (1, \dots, n)$, $\vec{u}_0 = (1, \dots, n+1)$, $\vec{v} = (n+3, \dots, n+m+2)$ and $\vec{v}_0 = (n+2, \dots, n+m+2)$.

The proof will be done by induction on $n+m$. For $n=m=0$, (2) implies

$$\varphi(XY) = \varphi(X)\varphi(Y) = b^1(X)\varphi(Y)$$

also, (4) gives

$$\varphi(XY) = b^2(XY) + b^1(X)\varphi(Y),$$

therefore $b^2(XY) = 0$.

Suppose the assertion is true for $n+m \leq N$ and let us prove it for $n+m = N$. From (4) we have that

$$\begin{aligned} \varphi(a_{\vec{w}}) &= \sum_{\vec{w}=\vec{w}_1, \vec{w}_2} b^{|\vec{w}_1|}(a_{\vec{w}_1})\varphi(a_{\vec{w}_2}) \\ &= \sum_{\substack{\vec{w}=\vec{w}_1, \vec{w}_2 \\ |\vec{w}_1| \leq n+1}} b^{|\vec{w}_1|}(a_{\vec{w}_1})\varphi(a_{\vec{w}_2}) + \sum_{\substack{\vec{w}=\vec{w}_1, \vec{w}_2 \\ |\vec{w}_1| > n+1}} b^{|\vec{w}_1|}(a_{\vec{w}_1})\varphi(a_{\vec{w}_2}) \end{aligned}$$

On the other hand, from (3),

$$\begin{aligned} \varphi(a_{\vec{w}}) &= \varphi(a_{\vec{u}}XYa_{\vec{v}}) \\ &= \varphi(a_{\vec{u}}X)\varphi(Ya_{\vec{v}}) = \varphi(a_{\vec{u}_0})\varphi(a_{\vec{v}_0}) \\ &= \sum_{\vec{u}_0=\vec{u}_1, \vec{u}_2} b(a_{\vec{u}_1})\varphi(a_{\vec{u}_2})\varphi(a_{\vec{v}_0}) \end{aligned}$$

Since the last element in $a_{\vec{u}_1}$ is $X \in \mathcal{A}_1$ and the first element in $a_{\vec{v}_0}$ is $Y \in \mathcal{A}_2$, property (3) gives

$$\varphi(a_{\vec{u}_2})\varphi(a_{\vec{v}_0}) = \varphi(a_{\overrightarrow{u_2, v_0}}).$$

Therefore

$$\begin{aligned} \varphi(a_{\vec{w}}) &= \sum_{\vec{u}_0=\vec{u}_1, \vec{u}_2} b(a_{\vec{u}_1})\varphi(a_{\overrightarrow{u_2, v_0}}) \\ &= \sum_{\substack{\vec{w}=\vec{w}_1, \vec{w}_2 \\ |\vec{w}_1| \leq n+1}} b^{|\vec{w}_1|}(a_{\vec{w}_1})\varphi(a_{\vec{w}_2}) \end{aligned}$$

It follows that

$$\sum_{\substack{\vec{w}=\vec{w}_1, \vec{w}_2 \\ |\vec{w}_1| > n+1}} b^{|\vec{w}_1|}(a_{\vec{w}_1})\varphi(a_{\vec{w}_2}) = 0$$

And the induction hypothesis implies the conclusion. $\square$

Corollary 3.2. *If X, Y are boolean independent, then*

$$B_{X+Y}(z) = B_X(z) + B_Y(z).$$

Proof. 3.2 is implied by $b_{X+Y}^n = b_X^n + b_Y^n$, which is an immediate consequence of 3.1 and the multilinearity of the mappings b^n . $\square$

The next proposition investigates properties of boolean cumulants with scalars among their entries.

Proposition 3.3. *If $n \geq 1$ and $a_1, \dots, a_n \in \mathcal{A}$, then*

- (i) $b^{n+1}(1, a_1, \dots, a_n) = 0$
- (ii) $b^{n+1}(a_1, \dots, a_n, 1) = 0$
- (iii) $b^{n+1}(a_1, \dots, X_k, 1, X_{k+1}, \dots, a_n) = b^n(a_1, \dots, a_n)$.

Proof. (i): Let $\vec{u} = (1, \dots, n)$, $\vec{v} = (1, \dots, k)$, $\vec{w} = (k+1, \dots, n)$.

$$\varphi(1, a_{\vec{u}}) = b^1(1)\varphi(a_{\vec{u}}) + \sum_{\vec{u}=\vec{u}_1, \vec{u}_2} b^{|\vec{u}_1|+1}(1, a_{\vec{u}_1})\varphi(a_{\vec{u}_2})$$

But $\varphi(1a_{\vec{u}}) = \varphi(a_{\vec{u}}) = b^1(1)\varphi(a_{\vec{u}})$, therefore

$$\sum_{\vec{u}=\vec{u}_1, \vec{u}_2} b^{|\vec{u}_1|+1}(1, a_{\vec{u}_1})\varphi(a_{\vec{u}_2}) = 0.$$

Since $\varphi(1X) = b^2(1, X) + b^1(1)\varphi(X)$, so $b^2(1, X) = 0$, property (ii) is proved by induction on n . The proof of (ii) is similar.

(iii): From (i), one has that

$$\begin{aligned} \varphi(a_1 1 a_2) &= b^3(a-1, 1, a_2) + b^2(a_1, 1)\varphi(a_2) + b^1(a_1)\varphi(1a_2) \\ &= b^3(a_1, 1, a_2) + b^1(a_1)\varphi(a_2). \end{aligned}$$

And, since $\varphi(a_1 a_2) = b^2(a_1, a_2) + b^1(a_1)\varphi(a_2)$ we have that $b^3((a_1, 1, a_2) = b^2(a_1, a_2)$.

The rest of the proof is again a induction on n :

$$\begin{aligned} \varphi(a_{\vec{v}} 1 a_{\vec{w}}) &= \sum_{\vec{v}=\vec{v}_1, \vec{v}_2} b^{|\vec{v}_1|}(a_{\vec{v}_1})\varphi(a_{\vec{v}_2} 1 a_{\vec{w}}) + b^{|\vec{v}|+1}\varphi(a_{\vec{w}}) + \\ &\quad \sum_{\vec{w}=\vec{w}_1, \vec{w}_2} b^{k+1+|\vec{w}_1|}(a_{\vec{u}}, 1, a_{\vec{w}_1})\varphi(a_{\vec{w}_2}) + b^{n+1}(a_{\vec{v}}, 1, a_{\vec{w}}) \end{aligned}$$

Applying (ii) and the induction hypothesis, the previous equation becomes

$$\begin{aligned} \varphi(a_{\vec{v}} 1 a_{\vec{w}}) &= \sum_{\vec{v}=\vec{v}_1, \vec{v}_2} b^{|\vec{v}_1|}(a_{\vec{v}_1})\varphi(a_{\vec{v}_2} 1 a_{\vec{w}}) + \sum_{\vec{w}=\vec{w}_1, \vec{w}_2} b^{k+|\vec{w}_1|}(a_{\vec{u}}, a_{\vec{w}_1})\varphi(a_{\vec{w}_2}) \\ &\quad + b^{n+1}(a_{\vec{v}}, 1, a_{\vec{w}}) \end{aligned}$$

and (2) gives the conclusion. $\square$

Corollary 3.4. *For all $X \in \mathcal{A}, n \geq 2$, one has that*

$$b_{1+X}^1 = 1 + b_X^1 \text{ and } b_{X+1}^n = \sum_{k=0}^{n-2} \binom{n-2}{k} b_X^{k+2}.$$

Proof. The multilinearity of b^n implies that:

$$b_{(1+X)}^n = \sum_{a_j \in \{1, X \mid j=1, \dots, n\}} b^n(a_1, a_2, \dots, a_n)$$

Terms having 1 as first or last entry also cancel from 3.3(i) and 3.3(ii). The remaining terms can have the entry 1 only on the remaining $n-2$ positions left, and 3.3(iii) gives the stated result. $\square$

Finally we will direct our attention toward boolean cumulants with products among their entries. The multiplicative property of the boolean cumulants will appear as a straightforward consequence of the following proposition.

Proposition 3.5. *Let $\mathcal{A}_1$ and $\mathcal{A}_2$ be two boolean independent subalgebras of $\mathcal{A}$, $X \in \mathcal{A}_1, Y \in \mathcal{A}_2$ and $a_1, \dots, a_n \in \mathcal{A}_1 \vee \mathcal{A}_2$. Then, for all $1 \leq k \leq n$*

- (i) $b^{n+1}(a_1, \dots, a_k, XY, a_{k+1}, \dots, a_n) = b^{k+1}(a_1, \dots, a_k, X)b^{n-k+1}(Y, a_{k+1}, \dots, a_n)$
- (ii) $b^{n+2}(a_1, \dots, a_k, XY, XY, a_{k+1}, \dots, a_n) = 0.$
- (iii) $b^{n+2}(a_1, \dots, a_k, Y, XY, a_{k+1}, \dots, a_n) = 0.$
- (iv) $b^{n+2}(a_1, \dots, a_k, XY, X, a_{k+1}, \dots, a_n) = 0.$

Proof. Again we use the short-hand notations $\vec{u} = (1, \dots, k), \vec{v} = (k+1, \dots, n)$.
 (i): For $n=0$, $\varphi(XY) = b^1(XY)$; also $\varphi(XY) = \varphi(X \cdot Y) = b^2(X, Y) + b^1(X)b^1(Y)$.
 But $b^2(X, Y) = 0$ from 3.1, hence $b^1(XY) = b^1(X)b^1(Y)$.

Suppose now 3.5(i) true for $n \leq N$ and let us prove it for $n = N+1$. One has that

$$\begin{aligned} \varphi(a_{\vec{u}}XYa_{\vec{v}}) &= \varphi(a_{\vec{u}}X)\varphi(Ya_{\vec{v}}) \\ &= \sum_{\vec{u}=\vec{u}_1, \vec{u}_2} b^{|\vec{u}_1|}(a_{\vec{u}_1})\varphi(a_{\vec{u}_2})\varphi(Ya_{\vec{v}}) + b^{k+1}(a_{\vec{u}}, X)\varphi(Ya_{\vec{v}}) \\ &= \sum_{\vec{u}=\vec{u}_1, \vec{u}_2} b^{|\vec{u}_1|}(a_{\vec{u}_1})\varphi(a_{\vec{u}_2})\varphi(Ya_{\vec{v}}) + b^{k+1}(a_{\vec{u}}, X)b^1(Y)\varphi(a_{\vec{v}}) \\ &\quad + \sum_{\vec{v}=\vec{v}_1, \vec{v}_2} b^{k+1}(a_{\vec{u}}, X)b^{|\vec{v}_1|+1}(Y, a_{\vec{v}_1})\varphi(a_{\vec{v}_2}) \\ &\quad + b^{k+1}(a_{\vec{u}}, X)b^{n-k+1}(Y, a_{\vec{v}}). \end{aligned}$$

Applying the induction hypothesis, the previous equality becomes

$$\begin{aligned} \varphi(a_{\vec{u}}XYa_{\vec{v}}) &= \sum_{\vec{u}=\vec{u}_1, \vec{u}_2} b^{|\vec{u}_1|}(a_{\vec{u}_1})\varphi(a_{\vec{u}_2})\varphi(Ya_{\vec{v}}) + b^{k+1}(a_{\vec{u}}, XY)\varphi(a_{\vec{v}}) \\ &\quad + \sum_{\vec{v}=\vec{v}_1, \vec{v}_2} b^{k+|\vec{v}_1|+1}(a_{\vec{u}}, XY, a_{\vec{v}_1})\varphi(a_{\vec{v}_2}) \\ &\quad + b^{k+1}(a_{\vec{u}}, X)b^{n-k+1}(Y, a_{\vec{v}}). \end{aligned}$$

Finally, comparing the above relation with the decomposition (3) for $\varphi(a_{\vec{u}} \cdot XY \cdot a_{\vec{v}})$, we obtain

$$b^{n+1}(a_{\vec{u}}, XY, a_{\vec{v}}) = b^{k+1}(a_{\vec{u}}, X)b^{n-k+1}(Y, a_{\vec{v}}).$$

(ii):Part (i) gives

$$\begin{aligned} b^{|\vec{u}|+|\vec{v}|+2}(a_{\vec{u}}, XY, XY, a_{\vec{v}}) &= b^{|\vec{u}|+1}(a_{\vec{u}}, X) \cdot b^{|\vec{v}|+2}(Y, XY, a_{\vec{v}}) \\ &= b^{|\vec{u}|+1}(a_{\vec{u}}, X) \cdot b^2(Y, X) \cdot b^{|\vec{v}|+1}(Y, a_{\vec{v}}) \\ &= 0. \end{aligned}$$

(iii) and (iv) are analogous. $\square$

Corollary 3.6. *For $\mathcal{A}_1, \mathcal{A}_2$ boolean independent algebras, $X \in \mathcal{A}_1, Y \in \mathcal{A}_2$ one has that:*

$$b_{X+Y+XY}^n = b_X^n + b_Y^n + \sum_{k=1}^n b_X^k b_Y^{n-k+1}$$

Proof. Let $\vec{u} = (1, \dots, n)$. From the multilinearity of b^n ,

$$b_{X+Y+XY}^n = \sum_{a_j \in \{X, Y, XY\}} b^n(a_{\vec{u}})$$

From 3.1, all the boolean cumulants containing both X and Y , but no XY vanish. From 3.5(ii)-(iv), among the boolean cumulants containing XY , only ones of the form $b^n(X, \dots, X, XY, Y, \dots, Y)$ are nonzero. Therefore:

$$b_{X+Y+XY}^n = b_X^n + b_Y^n + \sum_{k=0}^{n-1} b^n(\underbrace{X, \dots, X}_{k \text{ times}}, XY, Y, \dots, Y)$$

and the conclusion follows immediately from 3.5(i). $\square$

Corollary 3.7. *If X, Y are boolean independent, then*

$$(5) \quad B_{(1+X)(1+Y)}(z) = B_{(1+X)}(z) \cdot B_{(1+Y)}(z).$$

Proof. Let $Z = X + Y + XY$ and let α_n , respectively β_n be the coefficients of z^{n+1} from the left hand side, respectively the right hand side on 5. For $n = 1$,

$$\begin{aligned} \alpha_1 &= b_{(1+X)(1+Y)}^1 \\ &= 1 + b_X^1 + b_Y^1 + b_{XY}^1 \\ &= 1 + b_X^1 + b_Y^1 + b_X^1 b_Y^1 \\ &= (1 + b_X^1)(1 + b_Y^1) = b_{(1+X)}^1 b_{(1+Y)}^1 \\ &= \beta_1 \end{aligned}$$

For $n = 2$, we can write $\alpha_n = \sum_{\substack{p,q \geq 0 \\ p+q \geq 1}} \alpha_{p,q} b_X^p b_Y^q$ and $\beta_n = \sum_{\substack{p,q \geq 0 \\ p+q \geq 1}} \beta_{p,q} b_X^p b_Y^q$ and (5)

reduces to the equality between $\alpha_{p,q}$ and $\beta_{p,q}$.

Utilizing 3.4,

$$\begin{aligned} \alpha_n &= b_{1+Z}^n \\ &= \sum_{k=2}^n \binom{n-2}{k-2} b_Z^k \\ &= \sum_{k=2}^n \binom{n-2}{k-2} \left(b_X^k + b_Y^k + \sum_{l=1}^k b_X^l b_Y^{k-l+1} \right) \end{aligned}$$

Therefore

$$\alpha_{p,q} = \begin{cases} \binom{n-2}{p+q-2} & \text{if } p+q = n \text{ and } p=0 \text{ or } q=0 \\ \binom{n-2}{p+q-3} & \text{if } p, q \geq 1 \text{ and } p+q \leq n+1 \\ 0 & \text{otherwise} \end{cases}$$

On the other hand,

$$\begin{aligned} \beta_n &= \sum_{k=1}^n b_{(1+X)}^k b_{(1+Y)}^{n-k+1} \\ &= b_{(1+X)}^1 b_{(1+Y)}^n + b_{(1+X)}^n b_{(1+Y)}^1 \\ &\quad + \sum_{k=2}^{n-1} \left(\sum_{l=2}^k \binom{k-2}{l-2} b_X^l \sum_{s=2}^{n-k+1} \binom{n-k-1}{s-2} b_Y^s \right) \end{aligned}$$

Utilizing again 3.4 and that $b_{1+X}^1 = 1 + b_X^1$, we have that

$$\begin{aligned} \beta_n &= \sum_{k=2}^n \binom{n-2}{k-2} [b_X^k + b_Y^k + b_X^1 b_Y^k + b_X^k b_Y^1] \\ &\quad + \sum_{k=2}^{n-1} \left(\sum_{l=2}^k \sum_{s=2}^{n-k+1} \binom{k-2}{l-2} \binom{n-k-1}{s-2} b_X^l b_Y^s \right) \end{aligned}$$

therefore

$$\beta_{p,q} = \begin{cases} \binom{n-2}{p+q-2} & \text{if } p+q = n \text{ and } p=0 \text{ or } q=0 \\ \binom{n-2}{p+q-3} & \text{if } p=1 \text{ or } q=1 \text{ and } p+q \leq n+1 \\ \sum_{k=p}^{n-q+1} \binom{k-2}{s-2} \binom{n-k-1}{q-2} & \text{if } p, q \geq 2, \text{ and } p+q \leq n+1 \\ 0 & \text{otherwise} \end{cases}$$

The property (5) reduces though the to the equality

$$\sum_{k=p}^{n-q+1} \binom{k-2}{s-2} \binom{n-k-1}{q-2} = \binom{n-2}{p+q-3}$$

which is just an avatar of the well-known identity (see, for example, [7], Chapter 5):

$$\sum_{k=a}^{n-b} \binom{k}{a} \binom{n-k}{b} = \binom{n+1}{a+b+1}$$

□

4. BOOLEAN INDEPENDENCE WITH AMALGAMATION OVER AN ALGEBRA

4.1. Preliminaries. We need to consider an extended notion of non-unital complex algebra. $\mathfrak{A}$ will be called a $\mathfrak{B}$ -algebra if $\mathfrak{A}$ is an algebra such that $\mathfrak{B}$ is a subalgebra of $\mathfrak{A}$ or there is an algebra $\tilde{\mathfrak{A}}$ containing $\mathfrak{B}$ as a subalgebra such that $\tilde{\mathfrak{A}} = \mathfrak{A} \sqcup \mathfrak{B}$. (The symbol $\sqcup$ stands for disjoint union).

Let now $\mathfrak{A}$ be an algebra containing the unital subalgebra $\mathfrak{B}$ and $\Phi : \mathfrak{A} \longrightarrow \mathfrak{B}$ be a conditional expectation.

Suppose that $\mathfrak{A}_1$ and $\mathfrak{A}_2$ are two $\mathfrak{B}$ -subalgebras of $\mathfrak{A}$. We say that $\mathfrak{A}_1$ and $\mathfrak{A}_2$ are boolean independent with amalgamation over $\mathfrak{B}$ (or just boolean independent over $\mathfrak{B}$) if

$$\varphi(X_1 Y_1 X_2 \cdots) = \varphi(X_1) \varphi(Y_1) \cdots$$

for all $X_1, X_2, \dots \in \mathfrak{A}_1$ and $Y_1, Y_2, \dots \in \mathfrak{A}_2$.

Let $X \in \mathfrak{A}$ (if $\mathfrak{A}$ is a $*$ -algebra, we also require X to be selfadjoint). If X is not commuting with $\mathfrak{B}$, the algebra generated by X and $\mathfrak{B}$ will not be spanned by $\{\Phi(X^n)\}_n$. In this framework, the natural analogues of the n -th moment of X is the multilinear function $\widetilde{m}_X^n : \mathfrak{B}^{n-1} \rightarrow \mathfrak{B}$, given by

$$\widetilde{m}_X^n(f_1, \dots, f_{n-1}) = \Phi(X f_1 X \cdots X f_{n-1} X)$$

The objects corresponding to power series are elements from $Mul[[\mathfrak{B}]]$, the set of multilinear function series over $\mathfrak{B}$ (see [4]). A multilinear function series over $\mathfrak{B}$ is a sequence $F = (F_0, F_1, \dots)$ such that $F_0 \in \mathfrak{B}$ and F_n is a multilinear function from $\mathfrak{B}^n$ to $\mathfrak{B}$, for $n \geq 1$. For $F, G \in Mul[[\mathfrak{B}]]$, the *sum* $F + G$ and the *formal product* FG are the elements from $Mul[[\mathfrak{B}]]$ defined by:

$$\begin{aligned} (F + G)_n(f_1, \dots, f_n) &= F_n(f_1, \dots, f_n) + G_n(f_1, \dots, f_n) \\ (FG)_n(f_1, \dots, f_n) &= \sum_{k=0}^n F_k(f_1, \dots, f_k) G_{n-k}(f_{k+1}, \dots, f_n) \end{aligned}$$

Definition 4.1. *The object corresponding to the n -th boolean cumulant b^n (as defined in Section 2) is the multilinear function $\widetilde{b}^n : \mathfrak{A}^n \times \mathfrak{B}^{n-1} \rightarrow \mathfrak{B}$ given by the recurrence*

$$(6) \quad \Phi(X_1 f_1 X_2 \cdots X_{n-1} f_{n-1} X_n) = \sum_{k=1}^n \widetilde{b}_{X_1, \dots, X_k}^k(f_1, \dots, f_{k-1}) f_k \Phi(X_{k+1} f_{k+1} \cdots f_{n-1} X_n)$$

for all $X_1, \dots, X_n \in \mathfrak{A}$ and all $f_1, \dots, f_{n-1} \in \mathfrak{B}$. The n -uple $(X_1, \dots, X_n)$ will be called the lower argument of $\widetilde{b}^n$, while $(f_1, \dots, f_{n-1})$ will be called the upper argument of $\widetilde{b}^n$.

If $X_1 = \cdots = X_n = X$, we will write $\widetilde{b}_X^n$ for $\widetilde{b}_{X_1, \dots, X_n}^n$.

We will denote by $\widetilde{M}_X$, respectively $\widetilde{B}$ the multilinear function series for the multilinear moments and boolean cumulants of $X \in \mathfrak{A}$. With this notation, the recurrence in the definition of $\widetilde{b}^n$ can be rewritten as

$$\widetilde{M}_X = \widetilde{B}_X(1 + I\widetilde{M}_X)$$

for I the identity function on $\mathfrak{B}$.

Remark 4.2. *Boolean independence over the algebra $\mathfrak{B}$ implies boolean independence with respect to the functional Φ , in the sense of Section 2.*

We denote by b^n the n -th boolean cumulant with respect to Φ in the sense of Section 2. Comparing the recurrences (3) and (6), we have that

$$(7) \quad b^n(X_1, \dots, X_n) = \widetilde{b}_{X_1, \dots, X_n}^n(1, \dots, 1).$$

Since, for all $f_0, h_1, \dots, f_{n-1}h_n \in \mathfrak{B}$,

$$f_0 \Phi(X_1 h_1 f_1 X_2 \cdots h_{n-1} f_{n-1} X_n) h_n = \Phi((f_0 X_1 h_1) 1 (f_1 X_2 h_2) 1 \cdots 1 (f_{n-1} X_n h_n))$$

the recurrence (6) also gives

$$(8) \quad f_0 [\tilde{b}_{X_1, \dots, X_n}^n (h_1 f_1, \dots, h_{n-1} f_{n-1})] h_n = b^n (f_0 X_1 h_1, \dots, f_{n-1} X_n h_n).$$

4.2. Main results.

Proposition 4.3. *Let $\mathfrak{A}_1$ and $\mathfrak{A}_2$ independent over $\mathfrak{B}$ and $\mathfrak{A}_1 \vee_{\mathfrak{B}} \mathfrak{A}_2$ be the sub-algebra of $\mathfrak{A}$ generated by $\mathfrak{A}_1, \mathfrak{A}_2$ and $\mathfrak{B}$. Suppose that $X \in \mathfrak{A}_1$, $Y \in \mathfrak{A}_2$ and $X_1 \dots X_n \in \mathfrak{A}_1 \vee_{\mathfrak{B}} \mathfrak{A}_2$. Then*

- (i) $\tilde{b}_{X_1, \dots, X_k, X, Y, X_{k+1}, \dots, X_n}^{n+2} = 0$
- (ii) for all $f \in \mathfrak{B}$, one has that $\tilde{b}_{f, X_1, \dots, X_n}^{n+1} = \tilde{b}_{X_1, \dots, X_n, f}^{n+1} = 0$, while

$$\tilde{b}_{f, X_1, \dots, X_k, f, X_{k+1}, \dots, X_n}^{n+1} (f_1, \dots, f_n) = \tilde{b}_{X_1, \dots, X_n}^n (f_1, \dots, f_k f f_{k+1}, \dots, f_n)$$
- (iii) $\tilde{b}_{X_1, \dots, X_k, X Y, X_{k+1}, \dots, X_n}^{n+1} (f_1, \dots, f_n)$

$$= \tilde{b}_{X_1, \dots, X_k, X}^{k+1} (f_1, \dots, f_k) \tilde{b}_{Y, X_{k+1}, \dots, X_n}^{n-k+1} (f_{k+1}, \dots, f_n)$$

Proof. The proposition is an immediate consequence of Remark 4.2 and Propositions 3.3 and 3.5.

For (i) we need to prove that

$$\tilde{b}_{X_1, \dots, X_k, X, Y, X_{k+1}, \dots, X_n}^{n+2} (f_1, \dots, f_{n+1}) = 0$$

for all $f_1, \dots, f_{n+1} \in \mathfrak{B}$. But (8) implies

$$\tilde{b}_{X_1, \dots, X_k, X, Y, X_{k+1}, \dots, X_n}^{n+2} (f_1, \dots, f_{n+1}) = b^{n+2} (X_1 f_1, \dots, X f_{k+1}, Y f_{k+2}, \dots, X_n)$$

Since $X f_{k+1} \in \mathfrak{A}_1$ and $Y f_{k+2} \in \mathfrak{A}_2$, while $X_1 f_1, \dots, X_{n-1} f_{n+1}, X_n \in \mathfrak{A}_1 \vee_{\mathfrak{B}} \mathfrak{A}_2$, Proposition 3.1 implies the conclusion.

The parts (ii) and (iii) are analogous consequences of 4.2 and 3.3(i), 3.3(ii), respectively 3.5(i). $\square$

For stating the next results, we need a brief discussion about interval partitions.

Definition 4.4. *An interval partition γ on the set $\{1, 2, \dots, n\}$ is a collection of disjoint subsets, $D_1, \dots, D_q$, called blocks, such that*

- (i) $\bigcup_k = 1^q D_k = \{1, \dots, n\}$
- (ii) if $i_1 < i_2$ and $k \in D_{i_1}$, $l \in D_{i_2}$, then $k < l$.

The set of all interval partitions on $\{1, \dots, n\}$ will be denoted by $I(n)$. The number of blocks of the interval partition γ will be denoted by $|\gamma|$. For $\gamma_1 \in I(n)$ and $\gamma \in I(m)$, we will write $\gamma_1 \oplus \gamma_2$ for the interval partition from $I(m+n)$ obtained by juxtaposing γ_1 and γ_2 .

If $\gamma \in I(n-1)$, $\gamma = (1, \dots, p_1)(p_1 + 1, \dots, p_2), \dots, (p_{k-1} + 1, \dots, n-1)$, and $(f_1, \dots, f_n)$ is a n -uple from $\mathfrak{B}$, and then $\pi_\gamma(f_1, \dots, f_n)$ will denote the q -uple $(f_1 \cdots f_{p_1}, f_{p_1+1} \cdots f_{p_2}, \dots, f_{p_{q-1}+1} \cdots f_n)$ of elements from $\mathfrak{B}$.

Proposition 4.5. *For all $X \in \mathfrak{A}$ and $n \geq 2$, one has that*

$$(9) \quad \tilde{b}_{(1+X)}^n (f_1, \dots, f_{n-1}) = \sum_{\gamma \in I(n-1)} \tilde{b}_X^{|\gamma|+1} (\pi_\gamma(f_1, \dots, f_{n-1}))$$

Proof. From the multilinearity of $\tilde{b}^n$,

$$(10) \quad \tilde{b}_{(1+X)}^n = \sum_{\substack{X_j \in \{1, X\} \\ j=1, \dots, n}} \tilde{b}_{X_1, \dots, X_n}^n = \sum_{\substack{X_j \in \{1, X\} \\ j=2, \dots, n-1}} \tilde{b}_{X, X_2, \dots, X_{n-1}, X}^n,$$

since, from 4.3(ii), the terms with 1 on first and last position vanish.

Let $\gamma \in \mathcal{I}(n-1)$, $\gamma = (1, \dots, p_1)(p_1+1, \dots, p_2), \dots, (p_{k-1}+1, \dots, n-1)$. Denote by $\tilde{b}_{\gamma, X}^n$ the boolean cumulant having the lower argument $\underbrace{X, 1, \dots, 1}_{p_1}, \underbrace{X, 1, \dots, 1}_{p_2-p_1}, \dots, \underbrace{X, 1, \dots, 1}_{n-1-p_{k-1}}, X$. Then (10) becomes

$$\tilde{b}_{(1+X)}^n(f_1, \dots, f_{n-1}) = \sum_{\gamma \in \mathcal{I}(n-1)} \tilde{b}_{\gamma, X}^n(f_1, \dots, f_{n-1})$$

Finally, applying 4.3(iii), we get that

$$\begin{aligned} \tilde{b}_{\gamma, X}^n(f_1, \dots, f_{n-1}) &= \tilde{b}_X^{|\gamma|+1}(f_1 f_2 \cdots f_{p_1}, f_{p_1+1} \cdots f_{p_2}, \dots, f_{p_{k-1}+1} \cdots f_{n-1}) \\ &= \tilde{b}_X^{|\gamma|+1}(\pi_\gamma(f_1, \dots, f_n)), \end{aligned}$$

so q.e.d. $\square$

Corollary 4.6. *Suppose $\mathfrak{A}_1, \mathfrak{A}_2$ are boolean independent over $\mathfrak{B}$ and $X \in \mathfrak{A}_1$, $Y \in \mathfrak{A}_2$. Then*

$$\begin{aligned} \tilde{B}_{X+Y} &= \tilde{B}_X + \tilde{B}_Y \\ \tilde{B}_{(1+X)(1+Y)} &= \tilde{B}_{(1+X)} \tilde{B}_{(1+Y)}. \end{aligned}$$

Proof. The first relation amounts to showing that $\tilde{b}_{X+Y}^n = \tilde{b}_X^n + \tilde{b}_Y^n$, which is an obvious consequence of Proposition 4.3(i) and of the multilinearity of $\tilde{b}^n$.

For the second relation, first note that the n -th component of $B_{(1=x)(1+Y)}$ is

$$\begin{aligned} \tilde{b}_{(1+X)(1+Y)}^n(f_1, \dots, f_{n-1}) &= \tilde{b}_{1+(X+Y+XY)}^n(f_1, \dots, f_{n-1}) \\ &= \sum_{\gamma \in \mathcal{I}(n-1)} \tilde{b}_{X+Y+XY}^{|\gamma|+1}(f_1, \dots, f_{n-1}) \end{aligned}$$

The results 4.3(i) and (iii) imply the analogue of Corollary 3.6:

$$\begin{aligned} \tilde{b}_{X+Y+XY}^m(f_1, \dots, f_{m-1}) &= \left(\tilde{b}_X^m(1 + \tilde{b}_Y^1) + (1 + \tilde{b}_X^1) \tilde{b}_Y^m \right) (f_1, \dots, f_{m-1}) \\ &\quad + \sum_{k=2}^{m-2} \tilde{b}_X^k(f_1, \dots, f_{k-1}) \tilde{b}_Y^{m-k}(f_k, \dots, f_{m-k-1}) \end{aligned}$$

Fix $\gamma \in \mathcal{I}(n-1)$. The above equation gives

$$\begin{aligned} \tilde{b}_{X+Y+XY}^{|\gamma|+1}(\pi_\gamma(f_1, \dots, f_{n-1})) &= \left(\tilde{b}_X^{|\gamma|+1}(1 + \tilde{b}_Y^1) + (1 + \tilde{b}_X^1) \tilde{b}_Y^{|\gamma|+1} \right) (\pi_\gamma(f_1, \dots, f_{n-1})) \\ &\quad + \sum_{\substack{\gamma_1 \oplus \gamma_2 = \gamma \\ \gamma_1 \in \mathcal{I}(p) \\ \gamma_2 \in \mathcal{I}(n-p-1)}} \tilde{b}_X^{|\gamma_1|+1}(\pi_{\gamma_1}(f_1, \dots, f_p)) \tilde{b}_Y^{|\gamma_2|+1}(\pi_{\gamma_2}(f_{p+1}, \dots, f_{n-1})) \end{aligned}$$

therefore

$$\begin{aligned}
\tilde{b}_{(1+X)(1+Y)}^n(f_1, \dots, f_{n-1}) &= \sum_{\gamma \in \mathcal{I}(n-1)} \tilde{b}_{\gamma, X+Y+XY}^{|\gamma|+1}(\pi_\gamma(f_1, \dots, f_{n-1})) \\
&= \sum_{\gamma \in \mathcal{I}(n-1)} \tilde{b}_{X+Y+XY}^{|\gamma|+1}(\pi_\gamma(f_1, \dots, f_{n-1})) \\
&= \sum_{\gamma \in I(n-1)} \left(\tilde{b}_X^{|\gamma|+1}(1 + \tilde{b}_Y^1) + (1 + \tilde{b}_X^1) \tilde{b}_Y^{|\gamma|+1} \right) (\pi_\gamma(f_1, \dots, f_{n-1})) \\
&\quad + \sum_{k=2}^{n-1} \left(\sum_{\gamma \in I(k-1)} \tilde{b}_X^{|\gamma|+1}(\pi_\gamma(f_1, \dots, f_{k-1})) \right) \left(\sum_{\sigma \in I(n-k)} \tilde{b}_Y^{|\sigma|+1}(\pi_\sigma(f_k, \dots, f_{n-k})) \right)
\end{aligned}$$

Utilizing Prop(9), the above equation amounts to

$$\tilde{b}_{(1+X)(1+Y)}^n(f_1, \dots, f_{n-1}) = \sum_{k=1}^n \tilde{b}_{(1+X)}^k(f_1, \dots, f_{k-1}) \tilde{b}_{(1+Y)}^{n-k+1}(f_k, \dots, f_{n-1})$$

hence the conclusion. $\square$

Acknowledgements. My research was partially supported by the Grant 2-CEx06-11-34 of the Romanian Government. I am thankful to Șerban Belinschi, who brought to my attention the problem addressed in 4.6, to Hari Bercovici for reference [2] and to Marek Bożejko for reference [11].

REFERENCES

- [1] M. Anshelevich. Appell polynomials and their relatives II. Boolean theory. arXiv:0712.4185v1
- [2] H. Bercovici. On boolean convolutions Operator theory 20, Theta Ser. Adv. Math., vol. 6, Theta, Bucharest, 2006, 7–13.
- [3] M. Bożejko, M. Leinert and R. Speicher. Convolution and Limit Theorems for Conditionally free Random Variables. Pac. J. Math. 175 (1996), 357-388
- [4] K. Dykema. Multilinear function series and transforms in Free Probability theory. Preprint, arXiv:math.OA/0504361 v2 5 Jun 2005
- [5] K. Dykema. On the S-transform over a Banach algebra Preprint, arXiv:math.OA/0501083 01/2005
- [6] U. Franz. Boolean convolution of probability measures on the unit circle. arXiv:math.FA/0403243, 03/2004
- [7] R. Graham, D. Knuth, O. Patashnik Concrete mathematics, A Foundation for Computer Science Addison-Wesley Publishing Co., 1994
- [8] R.L. Hudson. A quantum mechanical central limit theorem for anticommuting observables J. Appl. Prob. 10(1973), 502–509
- [9] A. Nica, R. Speicher. Lectures on the Combinatorics of the Free Probability. London mathematical Society Lecture Note Series 335, Cambridge University Press 2006
- [10] M. Popa. A combinatorial approach to monotonic independence over a C^* -algebra Preprint, arXiv: math.OA/0612570, 01/2007
- [11] M. Schurmann, A. Ben Ghorbal. Non-commutative notions of stochastic independence. Math. Proc. Cambridge Philos. Soc. 133 (2002), no. 3, 531–561
- [12] R. Speicher. Combinatorial Theory of the Free Product with amalgamation and Operator-Valued Free Probability Theory. Mem. AMS, Vol 132, No 627 (1998)
- [13] R. Speicher, R. Waroudi Boolean convolution. Fields Institute Communications, Vol. 12 (D. Voiculescu, ed.), AMS, 1997, pp. 267-279
- [14] R.H. Terwiel. Projection operator method applied to stochastic linear differential equations Physica 74 (1974), 248–265

- [15] W. von Waldenfels. An approach to the theory of pressure broadening of spectral lines. Lecture Notes in Mathematics, vol. 296, Springer, Heidelberg, 1973, 19–69

¹INDIANA UNIVERSITY AT BLOOMINGTON, DEPARTMENT OF MATHEMATICS, RAWLES HALL, 931 E 3RD ST, BLOOMINGTON, IN 47405
E-mail address: `mipopa@indiana.edu`

²INSTITUTE OF MATHEMATICS, ROMANIAN ACADEMY, P.O.Box 1-764, BUCHAREST, RO-70700, ROMANIA